

DOULEUR : QUELLES ASSOCIATIONS ?

LEGENDE : ■ Association utile ■ Association possible (à discuter) ■ Association non possible	MEOPA	PARACETAMOL	NEFOPAM	CODEINE	TRAMADOL	NALBUPHINE	MORPHINE	MORPHINOMIMETIQUES	AINS	CORTICOIDES	BENZODIAZEPINES	ANESTHESIQUES LOCAUX
MEOPA (Medimix®)												
PARACETAMOL (Perfalgan®, Dafalgan®, Doliprane®, Efferalgan®)												
NEFOPAM (Acupan®)												
CODEINE (Dicodin LP®, Codenfan®)												
TRAMADOL (Contramal®)												
NALBUPHINE (Nubain®)												
MORPHINE (Morphine®, Actiskénan®, Skénan®)						*						
MORPHINOMIMETIQUES (Durogésic®, Actiq®, Sophidone®, Oxycontin®)												
AINS : KETOPROFENE (Profenid®), NAPROXENE (Apranax®)												
CORTICOIDES : PREDNISOLONE (Solumédrol®), PREDNISONE (Cortencyl®)												
BENZODIAZEPINES : (Midazolam®)				**	**	**	**	**				
ANESTHESIQUES LOCAUX : Xylocaïne®, Emla®												
TECHNIQUES NON MEDICAMENTEUSES : Respiration contrôlée, Diversion, Sophrologie, Hypnose, Musicothérapie, Arthérapie, Froid, Chaud, Immobilisation, Neurostimulation transcutanée												



* Sauf dans le cas Nalbuphine → Morphine, si Nalbuphine insuffisante

** Prudence